

**Michael P. Krumholz, M.D.
111 East 80th Street Suite 1C
New York, NY 10075
Tel. (212)734-5533/Fax (212)717-1688**

PREPARATION FOR COLONOSCOPY WITH GoLYTELY™(3 DOSES)

YOUR EXAMINATION WILL BE PERFORMED AT:

**Manhattan Endoscopy Center
535 Fifth Avenue, 5th Floor (entrance on 44th Street between Madison & Fifth Avenues)
New York, NY 10017
Tel. (212) 682-2828
Please arrive 30 minutes prior to your scheduled appointment time.**

IMPORTANT: Please read these instructions carefully **at least one week before** your examination.

PURCHASE: The office will send the prescription for the preparation kit(s) to the pharmacy, please provide the office with a pharmacy phone number at least 2 days before your Colonoscopy appointment. Also, purchase your choice of **clear fluids** from your grocery store. These include water, tea (hot or iced without milk or cream), apple juice, white grape juice, sports drinks (e.g., Gatorade), bouillon, broth (without noodles), Jell-O, sorbet, Italian ice, and popsicles. A container of Tucks or wet wipes (optional) may also be purchased.

MANAGEMENT OF MEDICATIONS AND DIET:

IMPORTANT: If you take any “blood thinners”, such as Coumadin (warfarin), Xarelto, Pradaxa, Eliquis, Plavix or Effient, you must consult with your cardiologist, neurologist or primary care physician **BEFORE** discontinuing these medications as outlined below. **Many patients with a history of heart disease, stents, clotting disorders, and stroke, may be advised to CONTINUE these drugs. If your doctor advises you to continue any of these medications, or advises an alternative “blood thinner”, please inform the office. We will need a letter of clearance from the prescriber to confirm specified instructions and that you are stable to proceed. It may also be necessary for Dr. Krumholz to consult with your doctor.**

If you are diabetic, check with your prescribing physician about taking your diabetes medication the morning of your procedure. Call your prescribing physician if you have questions regarding other medications. Otherwise, you may take your daily medications with a sip of water the morning of your procedure. Please bring a list of all your medications.

Seven days before your procedure

- Stop Plavix unless otherwise instructed by your physician.

Five days before your procedure

- Stop Coumadin (warfarin) unless otherwise instructed by your physician.
- Stop iron supplements.
- Avoid foods containing seeds (e.g., berries, tomatoes) and high fiber containing foods (e.g. raw fruit and vegetables, grains, nuts, beans).

Other medications referred to above may also need to be stopped for several days as determined by your physician. We may also recommend that aspirin and other non-steroidal anti-inflammatory drugs (NSAIDS) such as ibuprofen (Advil, Motrin), and naproxen (Aleve) be discontinued for several days depending on your clinical situation and the indication for your examination; if stopped, Tylenol (acetaminophen) may be substituted for pain relief.

One day before your procedure:

You will need to follow a **CLEAR FLUID DIET**– see list of examples above. You should not have **ANY** solid food again until after your colonoscopy. **IT IS VERY IMPORTANT THAT YOU DRINK PLENTY OF CLEAR FLUIDS, INCLUDING WATER, THROUGHOUT THE DAY. YOU SHOULD ATTEMPT TO CONSUME AT LEAST 2 LITERS OF FLUID.** Avoid juices with pulp such as orange or grapefruit juice. Plain teas are OK, but avoid dairy products, cream, and soy. You may use sugar, honey, or lemon. Avoid liquids that are red and blue. Please avoid alcohol.

- **Step 1: Mix the laxative preparation as directed on the label and place in the refrigerator to chill before drinking.**

- **Step 2: CONSUMING GOLYTELY™ SOLUTION**
 - **At 10:00am the DAY before your procedure,** Drink one 8 ounce glass every 15 minutes until **2 liters** is consumed. Drink the preparation with a straw; this may make it easier to consume. You may add the flavor packet provided with the medication, but **do not** add any additional flavoring to the liquid.
 - Drink 16 ounces of a clear liquid of your choice. You may start going to the bathroom after only a couple of glasses, but be sure to drink **ALL** of the solution. Refrigerate to chill the next dose of GOLYTELY™.

- **At 5:00pm the afternoon before your procedure repeat Step 2**

You may continue to drink clear fluids until bedtime. It is very important that you complete the entire laxative prep to ensure that the colon is cleansed. If you feel nauseated or bloated, stop drinking for a short while. This is usually temporary. Then try to continue drinking at a slower pace.

Usually liquid stools begin within two hours after you have taken the prep, though this may vary. Weakness may occur, especially if you have not taken enough fluid with the prep, so it is important to stay hydrated with other clear fluids.

On the day of your procedure:

- **Five hours before your scheduled procedure, repeat Step 2.** You may continue to drink clear fluids only until three hours before your examination.

Special Instructions:

- **Do not drink anything (except water with medications) for three hours before your examination**
- You will be sedated for the procedure so you must have an adult escort you home
- You will not be able to drive until the next morning
- Plan on being at Manhattan Endoscopy for approximately two hours
- Read all of the instructions and literature provided to you carefully
- Please inform the office if you have had an inadequate bowel preparation at the time of a previous examination or a history of liver or kidney failure

- **If you need to cancel or reschedule your appointment, please provide the office with three business days notice**
- Call the office if you have any questions or concerns